

PACO15679

Product Information

Size:

50ul

Reactivity:

Human, Mouse, Rat

Source:

Rabbit

Isotype:

IgG

Applications:

ELISA, WB, IHC

Recommended dilutions:

ELISA:1:1000-1:2000, WB:1:200-1:1000,
IHC:1:50-1:200

Protein Background:

STK40 (Serine/threonine-protein kinase 40), also known as Sugen kinase 495, is a 435 amino acid, protein kinase that may be a negative regulator of NF κ B and p53-mediated gene expression. Localized to both the cytoplasm and the nucleus, STK40 is ubiquitously expressed. STK40 has a central serine/threonine protein kinase domain and is homologous to TRB-3, a protein that regulates activation of MAP kinases and inhibits NF κ B-mediated gene transcription.

Gene ID:

STK40

Uniprot

Q8N2I9

Synonyms:

serine/threonine kinase 40


Immunogen:

Fusion protein of human STK40.

Storage:

-20 $\text{\textcircled{C}}$; C, pH7.4 PBS, 0.05% NaN₃, 40% Glycerol


Product Images



The image on the left is immunohistochemistry of paraffin-embedded Human thyroid cancer tissue using PACO15679(STK40 Antibody) at dilution 1/30, on the right is treated with fusion protein. (Original magnification: x—200).



Gel: 6%SDS-PAGE, Lysate: 40 μ g, Lane: Human placenta tissue, Primary antibody: PACO15679(STK40 Antibody) at dilution 1/400, Secondary antibody: Goat anti rabbit IgG at 1/8000 dilution, Exposure time: 2 minutes.



The image on the left is immunohistochemistry of paraffin-embedded Human cervical cancer tissue using PACO15679(STK40 Antibody) at dilution 1/30, on the right is treated with fusion protein. (Original magnification: x—200).