

### Product Information

**Size:**

50ul

**Reactivity:**

Human, Mouse, Rat

**Source:**

Rabbit

**Isotype:**

IgG

**Applications:**

ELISA, IHC

**Recommended dilutions:**

ELISA:1:2000-1:5000, IHC:1:50-1:200

**Protein Background:**

Phosphatidylinositol-4-phosphate-5-kinase (PIP5K) synthesizes phosphatidylinositol-4,5-bisphosphate, which regulates various processes including cell proliferation, survival, membrane trafficking, and cytoskeletal organization. The PIP5K family is divided into type I, type II and type III. Each type of the PIP5K family phosphorylate distinct substrates and they contain an activation loop, which determines their enzymatic specificity and subcellular targeting. The phosphatidylinositol-4-phosphate-5-kinase type I consists of three members, PIP5K I, integral, and , which are characterized by phosphorylating PI4P on the 5-hydroxyl. PIP5K I (designated PIP5K I integral in mouse) is expressed in brain tissue. PIP5K I integral, designated PIP5K I a in mouse, is also called STM7. PIP5K I has two variants produced by alternative splicing which are expressed in lung, brain, and kidneys.

**Gene ID:**

PIP5K1B

**Uniprot**

O14986

**Synonyms:**

phosphatidylinositol-4-phosphate 5-kinase, type I, beta

**Immunogen:**


Fusion protein of human PIP5K1B.

**Storage:**


-20&deg; C, pH7.4 PBS, 0.05% NaN<sub>3</sub>, 40% Glycerol

## Product Images

---



The image on the left is immunohistochemistry of paraffin-embedded Human liver cancer tissue using PACO15764(PIP5K1B Antibody) at dilution 1/30, on the right is treated with fusion protein. (Original magnification: x—200).



The image on the left is immunohistochemistry of paraffin-embedded Human thyroid cancer tissue using PACO15764(PIP5K1B Antibody) at dilution 1/30, on the right is treated with fusion protein. (Original magnification: x—200).