CAST Antibody



PACO15910

Reactivity:

Product Information

Size: Protein Background:

50ul The protein encoded by this gene is an endogenous calpain (calcium-dependent cysteine protease) inhibitor. It consists of an N-terminal domain L and four repetitive

calpain-inhibition domains (domains 1-4), and it is involved in the proteolysis of amyloid precursor protein. The calpain/calpastatin system is involved in numerous

Human amyloid precursor protein. The calpain/calpastatin system is involved in numerous membrane fusion events, such as neural vesicle exocytosis and platelet and red-cell aggregation. The encoded protein is also thought to affect the expression levels of

Rabbit genes encoding structural or regulatory proteins. Alternatively spliced transcript variants

encoding different isoforms have been described.

Isotype: Gene ID:

IgG CAST

Applications: Uniprot

ELISA, WB, IHC P20810

Recommended dilutions: Synonyms:

ELISA:1:2000-1:5000, WB:1:500-1:2000, calpastatin IHC:1:50-1:200


Immunogen:

Fusion protein of human CAST.


Storage:

-20° C, pH7.4 PBS, 0.05% NaN3, 40% Glycerol


Product Images



The image on the left is immunohistochemistry of paraffin-embedded Human colon cancer tissue using PACO15910(CAST Antibody) at dilution 1/60, on the right is treated with fusion protein. (Original magnification: x—200).



Gel: 8%SDS-PAGE, Lysate: 40 μ g, Lane: Lovo cells, Primary antibody: PACO15910(CAST Antibody) at dilution 1/600, Secondary antibody: Goat anti rabbit IgG at 1/8000 dilution, Exposure time: 7 seconds.



The image on the left is immunohistochemistry of paraffin-embedded Human cervical cancer tissue using PACO15910(CAST Antibody) at dilution 1/60, on the right is treated with fusion protein. (Original magnification: x—200).