

CRELD2 Antibody

PACO16072



Product Information	
Size:	Protein Background:
50ul	The epidermal growth factor (EGF) repeat-containing proteins constitute an expanding family of proteins that are involved in several cellular activities, such as blood coagulation, fibrinolysis, cell adhesion and neural and vertebrate development. CRELD2 (cysteine-rich with EGF-like domains 2) is a 353 amino acid, protein that is ubiquitously expressed and contains two FU domains and two EGF-like domains. Localized to the endoplasmic reticulum and secreted into the cell, CRELD2 interacts with AChR 4, possibly regulating its transport. Human CRELD2 shares 69% amino acid, identity with its mouse counterpart, suggesting a conserved role between species. Multiple isoforms of CRELD2 exist due to alternative splicing events. The gene encoding CRELD2 maps to human chromosome 22, which houses over 500 genes and is the second smallest human chromosome.
Reactivity:	
Human	
Source:	
Rabbit	
Isotype:	
lgG	
Applications:	Gene ID:
Elisa, WB, IHC	CRELD2
Recommended dilutions:	Uniprot
ELISA:1:2000-1:5000, WB:1:500-1:2000, IHC:1:50-1:200	Q6UXH1
	Synonyms:
	cysteine-rich with EGF-like domains 2
	Immunogen:
	Fusion protein of human CRELD2.
	Storage:
	-20° C, pH7.4 PBS, 0.05% NaN3, 40% Glycerol





The image on the left is immunohistochemistry of paraffin-embedded Human gastric cancer tissue using PACO16072(CRELD2 Antibody) at dilution 1/50, on the right is treated with fusion protein. (Original magnification: x—200).

Gel: 8%SDS-PAGE, Lysate: 40 μ g, Lane: Human normal colon tissue, Primary antibody: PACO16072(CRELD2 Antibody) at dilution 1/550, Secondary antibody: Goat anti rabbit IgG at 1/8000 dilution, Exposure time: 2 minutes.

The image on the left is immunohistochemistry of paraffin-embedded Human thyroid cancer tissue using PACO16072(CRELD2 Antibody) at dilution 1/50, on the right is treated with fusion protein. (Original magnification: x—200).