

HSD17B14 Antibody



PACO16493

Product Information

Size:

50ul

Reactivity:

Human, Mouse

Source:

Rabbit

Isotype:

IgG

Applications:

ELISA, WB, IHC

Recommended dilutions:

ELISA:1:1000-1:2000, WB:1:200-1:1000,
IHC:1:50-1:200

Protein Background:

17-beta-hydroxysteroid dehydrogenases, such as HSD17B14, are primarily involved in metabolism of steroids at the C17 position and also of other substrates, such as fatty acid, , prostaglandins, and xenobiotics.

Gene ID:

HSD17B14

Uniprot

Q9BPX1

Synonyms:

hydroxysteroid (17-beta) dehydrogenase 14


Immunogen:

Fusion protein of human HSD17B14.


Storage:

-20° C, pH7.4 PBS, 0.05% NaN₃, 40% Glycerol


Product Images



The image on the left is immunohistochemistry of paraffin-embedded Human liver cancer tissue using PACO16493(HSD17B14 Antibody) at dilution 1/30, on the right is treated with fusion protein. (Original magnification: x—200).



Gel: 10%SDS-PAGE, Lysate: 40 μ g, Lane: Mouse kidney tissue, Primary antibody: PACO16493(HSD17B14 Antibody) at dilution 1/200, Secondary antibody: Goat anti rabbit IgG at 1/8000 dilution, Exposure time: 3 minutes.



The image on the left is immunohistochemistry of paraffin-embedded Human colon cancer tissue using PACO16493(HSD17B14 Antibody) at dilution 1/30, on the right is treated with fusion protein. (Original magnification: x—200).