

Product Information

Size:

50ul

Reactivity:

Human, Mouse

Source:

Rabbit

Isotype:

IgG

Applications:

ELISA, WB, IHC

Recommended dilutions:

ELISA:1:2000-1:5000, WB:1:500-1:2000,
IHC:1:25-1:100

Protein Background:

The tetratricopeptide repeat (TPR) is a structural motif. It consists in a degenerate 34 amino acid, sequence motif identified in a wide variety of proteins. It is found in tandem arrays of 3–16 motifs, [1] which form scaffolds to mediate protein–protein interactions and often the assembly of multiprotein complexes. These repeats usually fold together to produce a single, linear solenoid domain called TPR domain.

Gene ID:

TTC23

Uniprot

Q5W5X9

Synonyms:

tetratricopeptide repeat domain 23


Immunogen:

Fusion protein of human TTC23.


Storage:

-20° C, pH7.4 PBS, 0.05% NaN₃, 40% Glycerol


Product Images



The image on the left is immunohistochemistry of paraffin-embedded Human liver cancer tissue using PACO17332(TTC23 Antibody) at dilution 1/30, on the right is treated with fusion protein. (Original magnification: x—200).



Gel: 8%SDS-PAGE, Lysate: 40 μ g, Lane 1-2: Mouse adrenal gland and kidney tissue, Primary antibody: PACO17332(TTC23 Antibody) at dilution 1/400 dilution, Secondary antibody: Goat anti rabbit IgG at 1/8000 dilution, Exposure time: 20 seconds.



The image on the left is immunohistochemistry of paraffin-embedded Human breast cancer tissue using PACO17332(TTC23 Antibody) at dilution 1/30, on the right is treated with fusion protein. (Original magnification: x—200).