## **THBS1 Antibody**

## PACO19219






Size:	Protein Background:
50ul	Non-classical major histocompatibility class lb molecule involved in immune self- nonself discrimination. In complex with B2M/beta-2-microglobulin binds nonamer self- peptides derived from the signal sequence of classical MHC class la molecules (VL9 peptides). Peptide-bound HLA-E-B2M heterotrimeric complex primarily functions as a ligand for natural killer (NK) cell inhibitory receptor KLRD1-KLRC1, enabling NK cells to monitor the expression of other MHC class I molecules in healthy cells and to tolerate self. Upon cellular stress, preferentially binds signal sequence-derived peptides from stress-induced chaperones and is no longer recognized by NK cell inhibitory receptor KLRD1-KLRC1, resulting in impaired protection from NK cells. Binds signal sequence- derived peptides from non-classical MHC class Ib HLA-G molecules and acts as a ligand for NK cell activating receptor KLRD1-KLRC2, likely playing a role in the generation and effector functions of adaptive NK cells and in maternal-fetal tolerance during pregnancy.
Reactivity:	
Human, Mouse	
Source:	
Rabbit	
lsotype:	
lgG	
Applications:	
Elisa, WB, IHC	Gene ID:
Recommended dilutions:	THBS1
ELISA:1:1000-1:2000, WB:1:200-1:1000, IHC:1:50-1:200	Uniprot
	P07996
	Synonyms:
	thrombospondin 1
	Immunogen:
	Synthetic peptide of human THBS1.

## Storage:

-20° C, pH7.4 PBS, 0.05% NaN3, 40% Glycerol



The image on the left is immunohistochemistry of paraffin-embedded Human cervical cancer tissue using PACO19219(THBS1 Antibody) at dilution 1/50, on the right is treated with synthetic peptide. (Original magnification: x—200).

Gel: 8%SDS-PAGE, Lysate: 40 μ g, Lane 1-2: A431 cells, Huvec cells, Primary antibody: PACO19219(THBS1 Antibody) at dilution 1/250, Secondary antibody: Goat anti rabbit IgG at 1/8000 dilution, Exposure time: 5 minutes.

The image on the left is immunohistochemistry of paraffin-embedded Human thyroid cancer tissue using PACO19219(THBS1 Antibody) at dilution 1/50, on the right is treated with synthetic peptide. (Original magnification: x—200).