KCNN4 Antibody



PACO19894

Product Information

Size:

50ul

Reactivity:

Human, Mouse

Source:

Rabbit

Isotype:

lgG

Applications:

ELISA, WB, IHC

ELISA:1:2000-1:5000, WB:1:500-1:2000,

Recommended dilutions:

IHC:1:50-1:200

Protein Background:

Important transcription factor regulating the expression of genes involved in immune and inflammatory responses. Plays also a significant role in adipogenesis, as well as in the gluconeogenic pathway, liver regeneration, and hematopoiesis. The consensus recognition site is 5'-T[TG]NNGNAA[TG]-3'. Its functional capacity is governed by protein interactions and post-translational protein modifications. During early embryogenesis, plays essential and redundant functions with CEBPA. Has a promitotic effect on many cell types such as hepatocytes and adipocytes but has an antiproliferative effect on T-cells by repressing MYC expression, facilitating differentiation along the T-helper 2 lineage. Binds to regulatory regions of several acute-phase and cytokines genes and plays a role in the regulation of acute-phase reaction and inflammation. Plays also a role in intracellular bacteria killing.

Gene ID:

KCNN4

Uniprot

O15554

Synonyms:

potassium intermediate/small conductance calcium-activated channel, subfamily N, member 4


Immunogen:

Synthetic peptide of human KCNN4.


Storage:

-20° C, pH7.4 PBS, 0.05% NaN3, 40% Glycerol


Product Images



The image on the left is immunohistochemistry of paraffin-embedded Human liver cancer tissue using PACO19894(KCNN4 Antibody) at dilution 1/40, on the right is treated with synthetic peptide. (Original magnification: x—200).



Gel: 8%SDS-PAGE, Lysate: 40 μ g, Lane: NIH/3T3 cells, Primary antibody: PACO19894(KCNN4 Antibody) at dilution 1/450, Secondary antibody: Goat anti rabbit IgG at 1/8000 dilution, Exposure time: 20 seconds.



The image on the left is immunohistochemistry of paraffin-embedded Human brain tissue using PACO19894(KCNN4 Antibody) at dilution 1/40, on the right is treated with synthetic peptide. (Original magnification: x—200).