

Product Information

Size:

50ul

Reactivity:

Human, Mouse

Source:

Rabbit

Isotype:

IgG

Applications:

ELISA, WB

Recommended dilutions:

ELISA:1:1000-1:2000, WB:1:200-1:1000

Protein Background:

Multifunctional protein that may modulate protein degradation pathways, apoptosis, transcription, signal transduction, cell cycle progress, and genetic stability by directly or indirectly interacting with hosts factors. Does not seem to be essential for HBV infection. May be directly involved in development of cirrhosis and liver cancer (hepatocellular carcinoma). Most of cytosolic activities involve modulation of cytosolic calcium. The effect on apoptosis is controversial depending on the cell types in which the studies have been conducted. By binding to human DDB1, may affect cell viability and stimulate genome replication. May induce apoptosis by localizing in mitochondria and causing loss of mitochondrial membrane potential. May also modulate apoptosis by binding human CFLAR, a key regulator of the death-inducing signaling complex (DISC). Moderately stimulates transcription of many different viral and cellular transcription elements.

Gene ID:

EMC1

Uniprot

Q8N766

Synonyms:

ER membrane protein complex subunit 1


Immunogen:

Synthetic peptide of human EMC1.

Storage:

-20° C, pH7.4 PBS, 0.05% NaN₃, 40% Glycerol

Product Images



Gel: 6%SDS-PAGE, Lysate: 40 μ g Primary antibody: PACO20272(EMC1 Antibody) at dilution 1/200 dilution, Secondary antibody: Goat anti rabbit IgG at 1/8000 dilution, Exposure time: 15 seconds.