## **EVA1A Antibody**


## PACO20713




Product Information	
Size:	Protein Background:
50ul	Component of the BRCA1-A complex, a complex that specifically recognizes 'Lys-63'-
Reactivity:	linked ubiquitinated histones H2A and H2AX at DNA lesions sites, leading to target the BRCA1-BARD1 heterodimer to sites of DNA damage at double-strand breaks (DSBs).
Human, Mouse	The BRCA1-A complex also possesses deubiquitinase activity that specifically removes 'Lys-63'-linked ubiquitin on histones H2A and H2AX. In the BRCA1-A complex, it acts as an adapter that bridges the interaction between BABAM1/NBA1 and the rest of the complex, thereby being required for the complex integrity and modulating the E3 ubiquitin ligase activity of the BRCA1-BARD1 heterodimer. Component of the BRISC complex, a multiprotein complex that specifically cleaves 'Lys-63'-linked ubiquitin in various substrates. Within the BRISC complex, acts as an adapter that bridges the interaction between BABAM1/NBA1 and the rest of the complex, thereby being required for the complex integrity. <b>Gene ID:</b> EVA1A <b>Uniprot</b>
Source:	
Rabbit	
lsotype:	
lgG	
Applications:	
ELISA, WB, IHC	
Recommended dilutions:	
ELISA:1:1000-1:2000, WB:1:200-1:1000,	
IHC:1:25-1:100	Q9H8M9
	Synonyms:
	eva-1 homolog A (C. elegans)
	Immunogen:
	Synthetic peptide of human EVA1A.
	<b>e</b> .

## Storage:

-20° C, pH7.4 PBS, 0.05% NaN3, 40% Glycerol





The image on the left is immunohistochemistry of paraffin-embedded Human thyroid cancer tissue using PACO20713(EVA1A Antibody) at dilution 1/30, on the right is treated with synthetic peptide. (Original magnification: x—200).

Gel: 12%SDS-PAGE, Lysate: 40 ug, Lane: Human normal liver tissue, Primary antibody: PACO20713(EVA1A Antibody) at dilution 1/550, Secondary antibody: Goat anti rabbit IgG at 1/8000 dilution, Exposure time: 10 seconds.

The image on the left is immunohistochemistry of paraffin-embedded Human liver cancer tissue using PACO20713(EVA1A Antibody) at dilution 1/30, on the right is treated with synthetic peptide. (Original magnification: x—200).