
VSIG10 Antibody

PACO20859



Product Information	
Size:	Protein Background:
50ul	Anti-apoptotic protein which acts by inhibiting the activities of CASP3, CASP7 and CASP9. Can inhibit the autocleavage of pro-CASP9 and cleavage of pro-CASP3 by CASP9. Capable of inhibiting CASP9 autoproteolysis at 'Asp-315' and decreasing the rate of auto proteolysis at 'Asp-330'. Acts as a mediator of neuronal survival in pathological conditions. Prevents motor-neuron apoptosis induced by a variety of signals. Possible role in the prevention of spinal muscular atrophy that seems to be caused by inappropriate persistence of motor-neuron apoptosis: mutated or deleted forms of NAIP have been found in individuals with severe spinal muscular atrophy. Acts as a sensor component of the NLRC4 inflammasome that specifically recognizes and binds needle protein Cprl from pathogenic bacteria C. violaceum. Association of pathogenic bacteria proteins drives in turn drive assembly and activation of the NLRC4 inflammasome, promoting caspase-1 activation, cytokine production and macrophage pyroptosis.
Reactivity:	
Human	
Source:	
Rabbit	
lsotype:	
lgG	
Applications:	
Elisa, ihc	Gene ID:
Recommended dilutions:	VSIG10
ELISA:1:2000-1:10000, IHC:1:30-1:150	Uniprot
	Q8N0Z9
	Synonyms:
	V-set and immunoglobulin domain containing 10
	Immunogen:
	Synthetic peptide of human VSIG10.
	Storage:

-20° C, pH7.4 PBS, 0.05% NaN3, 40% Glycerol



The image on the left is immunohistochemistry of paraffin-embedded Human thyroid cancer tissue using PACO20859(VSIG10 Antibody) at dilution 1/45, on the right is treated with synthetic peptide. (Original magnification: x—200).

The image on the left is immunohistochemistry of paraffin-embedded Human brain tissue using PACO20859(VSIG10 Antibody) at dilution 1/45, on the right is treated with synthetic peptide. (Original magnification: x—200).