

PACO20953

Product Information

Size:

50ul

Reactivity:

Human, Mouse, Rat

Source:

Rabbit

Isotype:

IgG

Applications:

ELISA, WB, IHC

Recommended dilutions:

ELISA:1:2000-1:5000, WB:1:500-1:2000,
IHC:1:20-1:100

Protein Background:

Secreted effector that interferes with the host cell ubiquitin pathway and is required for intracellular bacterial replication. Catalyzes the ubiquitination of several mammalian Rab proteins (Rab33b, Rab1, Rab6a and Rab30) during *L. pneumophila* infection, without engaging the standard cellular enzyme cascade (E1 and E2). Transfers an ADP-ribose moiety from NAD to the 'Arg-42' residue of ubiquitin in a reaction that releases nicotinamide. The modified ubiquitin is subsequently transferred to the substrate protein through an unknown mechanism that results in the release of AMP. Cannot ubiquitinate the endosomal Rab5 or the cytoskeletal small GTPase Rac1. Also acts as a deubiquitinase (DUB), catalyzing the cleavage of three of the most abundant polyubiquitin chains ('Lys-11', 'Lys-48' and 'Lys-63') with a distinct preference for 'Lys-63' linkages; is thus able to efficiently remove 'Lys-63'-linked polyubiquitin chains from the phagosomal surface. Is also able to remove NEDD8 from neddylated proteins, but is unable to recognize SUMO.

Gene ID:

ARFGAP2

Uniprot

Q8N6H7

Synonyms:

ADP-ribosylation factor GTPase activating protein 2


Immunogen:

Synthetic peptide of human ARFGAP2.


Storage:

-20° C, pH7.4 PBS, 0.05% NaN3, 40% Glycerol

Product Images



Gel: 8%SDS-PAGE, Lysate: 40 ug, Lane 1-2: A431 and hela cell, Primary antibody: PACO20953(ARFGAP2 Antibody) at dilution 1/200 dilution, Secondary antibody: Goat anti rabbit IgG at 1/8000 dilution, Exposure time: 15 seconds.



The image on the left is immunohistochemistry of paraffin-embedded Human colorectal cancer tissue using PACO20953(ARFGAP2 Antibody) at dilution 1/25, on the right is treated with synthetic peptide. (Original magnification: x—200).