P4HB Antibody, FITC conjugated

PACO25210



Product Information	
Size:	Protein Background:
50ug	Dsb proteins (DsbA, DsbB, DsbC, and DsbD) catalyze formation and isomerization of protein disulfide bonds in the periplasm of Escherichia coli. Disulphide bond isomerase (DsbC), a member of the thioredoxin superfamily, acts as a disulfide isomerase and converts aberrant disulfide bonds to correct ones. It also acts by transferring its disulfide bond to other proteins and is reduced in the process. DsbC is reoxidized by a yet uncharacterized protein.
Reactivity:	
Human	
Source:	
Rabbit	Gene ID:
lsotype:	Р4НВ
lgG	Uniprot
Applications:	P07237
ELISA	Synonyms:
Recommended dilutions:	Protein disulfide-isomerase (PDI) (EC 5.3.4.1) (Cellular thyroid hormone-binding protein) (Prolyl 4-hydroxylase subunit beta) (p55), P4HB, ERBA2L PDI PDIA1 PO4DB
	Immunogen:
	Recombinant Human Protein disulfide-isomerase protein (29-508AA).
	Storage:
	Preservative: 0.03% Proclin 300. Constituents: 50% Glycerol, 0.01M PBS, PH 7.4

N/A N/A