## P4HB Antibody, Biotin conjugated



## PACO25211

Reactivity:

Human

Source:

## **Product Information**

Size: Protein Background:

50ug Dsb proteins (DsbA, DsbB, DsbC, and DsbD) catalyze formation and isomerization of

protein disulfide bonds in the periplasm of Escherichia coli. Disulphide bond isomerase (DsbC), a member of the thioredoxin superfamily, acts as a disulfide isomerase and

converts aberrant disulfide bonds to correct ones. It also acts by transferring its disulfide

bond to other proteins and is reduced in the process. DsbC is reoxidized by a yet

uncharacterized protein.

Rabbit Gene ID:

**Isotype:** P4HB

lgG Uniprot

**Applications:** P07237

ELISA Synonyms:

Recommended dilutions: Protein disulfide-isomerase (PDI) (EC 5.3.4.1) (Cellular thyroid hormone-binding protein)

(Prolyl 4-hydroxylase subunit beta) (p55), P4HB, ERBA2L PDI PDIA1 PO4DB

Immunogen:

Recombinant Human Protein disulfide-isomerase protein (29-508AA).

Storage:

Preservative: 0.03% Proclin 300. Constituents: 50% Glycerol, 0.01M PBS, PH 7.4

Product	<b>Images</b>
---------	---------------

N/A N/A