FUOM Antibody, Biotin conjugated

PACO25303



| Product Information | |
|------------------------|---|
| Size: | Protein Background: |
| 50ug | Involved in the interconversion between alpha- and beta-L-fucoses. L-Fucose (6-deoxy- |
| Reactivity: | L-galactose) exists as alpha-L-fucose (29.5%) and beta-L-fucose (70.5%), the beta-form is metabolized through the salvage pathway. GDP-L-fucose formed either by the de |
| Human | novo or salvage pathways is transported into the endoplasmic reticulum, where it serves as a substrate for N- and O-glycosylations by fucosyltransferases. Fucosylated |
| Source: | structures expressed on cell surfaces or secreted in biological fluids are believed to play |
| Rabbit | a critical role in cell-cell adhesion and recognition processes. |
| lsotype: | Gene ID: |
| lgG | FUOM |
| Applications: | Uniprot |
| ELISA | A2VDF0 |
| Recommended dilutions: | Synonyms: |
| | Fucose mutarotase (EC 5.1.3.29), FUOM, C10orf125 |
| | Immunogen: |
| | Recombinant Human Fucose mutarotase protein (1-134AA). |
| | Storage: |
| | Preservative: 0.03% Proclin 300. Constituents: 50% Glycerol, 0.01M PBS, PH 7.4 |

N/A N/A