

KCNJ10 Antibody, FITC conjugated



PACO27755

Product Information

Size:

50ug

Reactivity:

Human

Source:

Rabbit

Isotype:

IgG

Applications:

ELISA

Recommended dilutions:

Protein Background:

May be responsible for potassium buffering action of glial cells in the brain. Inward rectifier potassium channels are characterized by a greater tendency to allow potassium to flow into the cell rather than out of it. Their voltage dependence is regulated by the concentration of extracellular potassium; as external potassium is raised, the voltage range of the channel opening shifts to more positive voltages. The inward rectification is mainly due to the blockage of outward current by internal magnesium. Can be blocked by extracellular barium and cesium.

Gene ID:

KCNJ10

Uniprot

P78508

Synonyms:

ATP-sensitive inward rectifier potassium channel 10 (ATP-dependent inwardly rectifying potassium channel Kir4.1) (Inward rectifier K(+) channel Kir1.2) (Potassium channel, inwardly rectifying subfamily J member 10), KCNJ10

Immunogen:

Recombinant Human ATP-sensitive inward rectifier potassium channel 10 protein (165-379AA).

Storage:

Preservative: 0.03% Proclin 300. Constituents: 50% Glycerol, 0.01M PBS, PH 7.4

Product Images

N/A

N/A