MED24 Antibody, Biotin conjugated

PACO28209



Product Information	
Size:	Protein Background:
50ug	Component of the Mediator complex, a coactivator involved in the regulated
Reactivity:	transcription of nearly all RNA polymerase II-dependent genes. Mediator functions as a bridge to convey information from gene-specific regulatory proteins to the basal RNA
Human	polymerase II transcription machinery. Mediator is recruited to promoters by direct interactions with regulatory proteins and serves as a scaffold for the assembly of a
Source:	functional preinitiation complex with RNA polymerase II and the general transcription
Rabbit	factors.
lsotype:	Gene ID:
lgG	MED24
Applications:	Uniprot
ELISA	O75448
Recommended dilutions:	Synonyms:
	Mediator of RNA polymerase II transcription subunit 24 (Activator-recruited cofactor 100 kDa component) (ARC100) (Cofactor required for Sp1 transcriptional activation subunit 4) (CRSP complex subunit 4) (Mediator complex subunit 24) (Thyroid hormone receptor-associated protein 4) (Thyroid hormone receptor-associated protein complex 100 kDa component)
	Immunogen:
	Recombinant Human Mediator of RNA polymerase II transcription subunit 24 protein (1-300AA).
	Storage:

Preservative: 0.03% Proclin 300. Constituents: 50% Glycerol, 0.01M PBS, PH 7.4

N/A N/A