
RBPMS Antibody

PACO31856



| Product Information | |
|--|--|
| Size: | Protein Background: |
| 50ug | Acts as a coactivator of transcriptional activity. Required to increase TGFB1/Smad- mediated transactivation. Acts through SMAD2, SMAD3 and SMAD4 to increase transcriptional activity. Increases phosphorylation of SMAD2 and SMAD3 on their C- terminal SSXS motif, possibly through recruitment of TGFBR1. Promotes the nuclear accumulation of SMAD2, SMAD3 and SMAD4 proteins. Binds to poly(A) RNA. Gene ID: |
| Reactivity: | |
| Human | |
| Source: | |
| Rabbit | RBPMS |
| lsotype: | Uniprot |
| lgG | Q93062 |
| Applications: | Synonyms: |
| ELISA, IHC, IF | RNA-binding protein with multiple splicing (RBP-MS) (Heart and RRM expressed |
| Recommended dilutions: | sequence) (Hermes), RBPMS, HERMES |
| ELISA:1:2000-1:10000, IHC:1:20-1:200, IF:1:50-1:200 | Immunogen: |
| | Recombinant Human RNA-binding protein with multiple splicing protein (1-219AA). |
| | Storage: |

Preservative: 0.03% Proclin 300. Constituents: 50% Glycerol, 0.01M PBS, PH 7.4



Immunohistochemistry of paraffin-embedded human kidney tissue using PACO31856 at dilution of 1:100.

Immunofluorescent analysis of PC-3 cells using PACO31856 at dilution of 1:100 and Alexa Fluor 488-congugated AffiniPure Goat Anti-Rabbit IgG(H+L).