

CCL14 Antibody, FITC conjugated



PACO32638

Product Information

Size:

50ug

Reactivity:

Human

Source:

Rabbit

Isotype:

IgG

Applications:

ELISA

Recommended dilutions:

Protein Background:

Has weak activities on human monocytes and acts via receptors that also recognize MIP-1 alpha. It induced intracellular Ca²⁺ changes and enzyme release, but no chemotaxis, at concentrations of 100-1,000 nM, and was inactive on T-lymphocytes, neutrophils, and eosinophil leukocytes. Enhances the proliferation of CD34 myeloid progenitor cells. The processed form HCC-1(9-74) is a chemotactic factor that attracts monocytes eosinophils, and T-cells and is a ligand for CCR1, CCR3 and CCR5.

Gene ID:

CCL14

Uniprot

Q16627

Synonyms:

C-C motif chemokine 14 (Chemokine CC-1/CC-3) (HCC-1/HCC-3) (HCC-1(1-74)) (NCC-2) (Small-inducible cytokine A14) [Cleaved into: HCC-1(3-74); HCC-1(4-74); HCC-1(9-74)], CCL14, NCC2 SCYA14

Immunogen:

Recombinant Human C-C motif chemokine 14 protein (20-93AA).

Storage:

Preservative: 0.03% Proclin 300. Constituents: 50% Glycerol, 0.01M PBS, PH 7.4

Product Images

N/A

N/A