S100B Antibody, Biotin conjugated

PACO32891



Product Information	
Size:	Protein Background:
50ug	Weakly binds calcium but binds zinc very tightly-distinct binding sites with different
Reactivity:	affinities exist for both ions on each monomer. Physiological concentrations of potassium ion antagonize the binding of both divalent cations, especially affecting
Human	high-affinity calcium-binding sites. Binds to and initiates the activation of STK38 by releasing autoinhibitory intramolecular interactions within the kinase. Interaction with
Source:	AGER after myocardial infarction may play a role in myocyte apoptosis by activating
Rabbit	ERK1/2 and p53/TP53 signaling. Could assist ATAD3A cytoplasmic processing, preventing aggregation and favoring mitochondrial localization. May mediate calcium-
lsotype:	dependent regulation on many physiological processes by interacting with other proteins, such as TPR-containing proteins, and modulating their activity.
lgG	Gene ID:
Applications:	S100B
ELISA	Uniprot
Recommended dilutions:	P04271
	Synonyms:
	Protein S100-B (S-100 protein beta chain) (S-100 protein subunit beta) (S100 calcium- binding protein B), S100B
	Immunogen:
	Recombinant Human Protein S100-B protein (2-92AA).
	Storage:

Preservative: 0.03% Proclin 300. Constituents: 50% Glycerol, 0.01M PBS, PH 7.4

N/A N/A