

Product Information

Size:

50ug

Reactivity:

Human

Source:

Rabbit

Isotype:

IgG

Applications:

ELISA, IHC

Recommended dilutions:

ELISA:1:2000-1:10000, IHC:1:20-1:200

Protein Background:

Binds and transactivates the retinoic acid, response elements that control expression of the retinoic acid, receptor beta 2 and alcohol dehydrogenase 3 genes. Transactivates both the phenobarbital responsive element module of the human CYP2B6 gene and the CYP3A4 xenobiotic response element.

Gene ID:

NR113

Uniprot

Q14994

Synonyms:

Nuclear receptor subfamily 1 group I member 3 (Constitutive activator of retinoid response) (Constitutive active response) (Constitutive androstane receptor) (CAR) (Orphan nuclear receptor MB67), NR113, CAR


Immunogen:

Recombinant Human Nuclear receptor subfamily 1 group I member 3 protein (2-351AA).


Storage:

Preservative: 0.03% Proclin 300. Constituents: 50% Glycerol, 0.01M PBS, PH 7.4


Product Images



Immunohistochemistry of paraffin-embedded human kidney tissue using PACO33292 at dilution of 1:100.



Immunohistochemistry of paraffin-embedded human liver tissue using PACO33292 at dilution of 1:100.



Immunohistochemistry of paraffin-embedded human testis tissue using PACO33292 at dilution of 1:100.