CXXC5 Antibody, HRP conjugated

PACO36739



Product Information	
Size:	Protein Background:
50ug	May indirectly participate in activation of the NF-kappa-B and MAPK pathways. Acts as
Reactivity:	a mediator of BMP4-mediated modulation of canonical Wnt signaling activity in neural stem cells. Required for DNA damage-induced ATM phosphorylation, p53 activation
Human	and cell cycle arrest. Involved in myelopoiesis. Transcription factor. Binds to the oxygen responsive element of COX4I2 and represses its transcription under hypoxia conditions (4% oxygen), as well as normoxia conditions (20% oxygen). May repress COX4I2 transactivation induced by CHCHD2 and RBPJ.
Source:	
Rabbit	
lsotype:	Gene ID:
IgG	CXXC5
Applications:	Uniprot
ELISA	Q7LFL8
Recommended dilutions:	Synonyms:
	CXXC-type zinc finger protein 5 (CF5) (Putative MAPK-activating protein PM08) (Putative NF-kappa-B-activating protein 102) (Retinoid-inducible nuclear factor) (RINF), CXXC5
	Immunogen:
	Recombinant Human CXXC-type zinc finger protein 5 protein (1-227AA).
	Storage:
	Preservative: 0.03% Proclin 300. Constituents: 50% Glycerol, 0.01M PBS, PH 7.4

N/A N/A