

CXXC5 Antibody, FITC conjugated



PACO36740

Product Information

Size:

50ug

Reactivity:

Human

Source:

Rabbit

Isotype:

IgG

Applications:

ELISA

Recommended dilutions:

Protein Background:

May indirectly participate in activation of the NF-kappa-B and MAPK pathways. Acts as a mediator of BMP4-mediated modulation of canonical Wnt signaling activity in neural stem cells. Required for DNA damage-induced ATM phosphorylation, p53 activation and cell cycle arrest. Involved in myelopoiesis. Transcription factor. Binds to the oxygen responsive element of COX4I2 and represses its transcription under hypoxia conditions (4% oxygen), as well as normoxia conditions (20% oxygen). May repress COX4I2 transactivation induced by CHCHD2 and RBPJ.

Gene ID:

CXXC5

Uniprot

Q7LFL8

Synonyms:

CXXC-type zinc finger protein 5 (CF5) (Putative MAPK-activating protein PM08) (Putative NF-kappa-B-activating protein 102) (Retinoid-inducible nuclear factor) (RINF), CXXC5

Immunogen:

Recombinant Human CXXC-type zinc finger protein 5 protein (1-227AA).

Storage:

Preservative: 0.03% Proclin 300. Constituents: 50% Glycerol, 0.01M PBS, PH 7.4

Product Images

N/A

N/A