

ATG4D Antibody, FITC conjugated



PACO37464

Product Information

Size:

50ug

Reactivity:

Human

Source:

Rabbit

Isotype:

IgG

Applications:

ELISA

Recommended dilutions:

Protein Background:

Cysteine protease ATG4D: Cysteine protease required for the cytoplasm to vacuole transport (Cvt) and autophagy. Cleaves the C-terminal amino acid, of ATG8 family proteins MAP1LC3 and GABARAPL2, to reveal a C-terminal glycine. Exposure of the glycine at the C-terminus is essential for ATG8 proteins conjugation to phosphatidylethanolamine (PE) and insertion to membranes, which is necessary for autophagy. Has also an activity of delipidating enzyme for the PE-conjugated forms.

Gene ID:

ATG4D

Uniprot

Q86TLO

Synonyms:

Cysteine protease ATG4D (EC 3.4.22. -) (AUT-like 4 cysteine endopeptidase) (Autophagin-4) (Autophagy-related cysteine endopeptidase 4) (Autophagy-related protein 4 homolog D) [Cleaved into: Cysteine protease ATG4D, mitochondrial], ATG4D, APG4D AUTL4

Immunogen:

Recombinant Human Cysteine protease ATG4D protein (14-43AA).

Storage:

Preservative: 0.03% Proclin 300. Constituents: 50% Glycerol, 0.01M PBS, PH 7.4

Product Images

N/A

N/A