## WDYHV1 Antibody, HRP conjugated

## PACO39595



Product Information	
Size:	Protein Background:
50ug	Mediates the side-chain deamidation of N-terminal glutamine residues to glutamate,
Reactivity:	an important step in N-end rule pathway of protein degradation. Conversion of the resulting N-terminal glutamine to glutamate renders the protein susceptible to
Human	arginylation, polyubiquitination and degradation as specified by the N-end rule. Does not act on substrates with internal or C-terminal glutamine and does not act on non-
Source:	glutamine residues in any position. Does not deaminate acetylated N-terminal
Rabbit	glutamine. With the exception of proline, all tested second-position residues on substrate peptides do not greatly influence the activity. In contrast, a proline at position
lsotype:	2, virtually abolishes deamidation of N-terminal glutamine.
lgG	Gene ID:
Applications:	WDYHV1
ELISA	Uniprot
Recommended dilutions:	Q96HA8
	Synonyms:
	Protein N-terminal glutamine amidohydrolase (EC 3.5.1.122) (Protein NH2-terminal glutamine deamidase) (N-terminal Gln amidase) (Nt(Q)-amidase) (WDYHV motif- containing protein 1), WDYHV1, C8orf32 NTAQ1
	Immunogen:
	Recombinant Human Protein N-terminal glutamine amidohydrolase protein (1-205AA).
	Storage:
	Preservative: 0.03% Proclin 300. Constituents: 50% Glycerol, 0.01M PBS, PH 7.4

N/A N/A