# **CIART Antibody**



### PACO39638

#### **Product Information**

Size:

50ug

Reactivity:

Human

Source: Rabbit

Isotype:

lgG

**Applications:** 

ELISA, IHC

Recommended dilutions:

ELISA:1:2000-1:10000, IHC:1:20-1:200

### **Protein Background:**

Transcriptional repressor which forms a negative regulatory component of the circadian clock and acts independently of the circadian transcriptional repressors: CRY1, CRY2 and BHLHE41. In a histone deacetylase-dependent manner represses the transcriptional activator activity of the CLOCK-ARNTL/BMAL1 heterodimer. Abrogates the interaction of ARNTL/BMAL1 with the transcriptional coactivator CREBBP and can repress the histone acetyl-transferase activity of the CLOCK-ARNTL/BMAL1 heterodimer, reducing histone acetylation of its target genes. Rhythmically binds the E-box elements (5'-CACGTG-3') on circadian gene promoters and its occupancy shows circadian oscillation antiphasic to ARNTL/BMAL1. Interacts with the glucocorticoid receptor (NR3C1) and contributes to the repressive function in the glucocorticoid response.

Gene ID:

CIART

Uniprot

Q8N365

## **Synonyms:**

Circadian-associated transcriptional repressor (ChIP-derived repressor of network oscillator) (Chrono) (Computationally highlighted repressor of the network oscillator), CIART, C1orf51


### Immunogen:

Recombinant Human Circadian-associated transcriptional repressor protein (1-385AA).

## Storage:

Preservative: 0.03% Proclin 300. Constituents: 50% Glycerol, 0.01M PBS, PH 7.4

## **Product Images**



Immunohistochemistry of paraffin-embedded human thyroid tissue using PACO39638 at dilution of 1:100.