## SPRTN Antibody, HRP conjugated



## PACO40535

## **Product Information**

Size: Protein Background:

50ug Regulator of UV-induced DNA damage response: acts as a 'reader' of ubiquitinated PCNA that enhances RAD18-mediated PCNA ubiquitination and translesion DNA

Reactivity: synthesis (TLS). Recruited to sites of UV damage and interacts with ubiquitinated PCNA and RAD18, the E3 ubiquitin ligase that monoubiquitinates PCNA. Facilitates chromatin

association of RAD18 and is required for efficient PCNA monoubiquitination, promoting

Source:

a feed-forward loop to enhance PCNA ubiquitination and translesion DNA synthesis

**ource:**a feed-forward loop to enhance PCNA ubiquitination and translesion DNA synthesis.
Acts as a regulator of TLS by recruiting VCP/p97 to sites of DNA damage, possibly

Rabbit leading to extraction of DNA polymerase eta (POLH) by VCP/p97 to prevent excessive

**Isotype:** translesion DNA synthesis and limit the incidence of mutations induced by DNA

damage.

lgG **Gene ID:** 

Applications:

ELISA Uniprot

Recommended dilutions: Q9H040

Synonyms:

SprT-like domain-containing protein Spartan (DNA damage protein targeting VCP) (DVC1) (Protein with SprT-like domain at the N terminus) (Spartan), SPRTN, C1orf124

DVC1

Immunogen:

Recombinant Human SprT-like domain-containing protein Spartan protein (1-239AA).

Storage:

Preservative: 0.03% Proclin 300. Constituents: 50% Glycerol, 0.01M PBS, PH 7.4

Product	<b>Images</b>
---------	---------------

N/A N/A