MED28 Antibody, Biotin conjugated



PACO40553

Rabbit

Product Information

Size: Protein Background:

50ug Component of the Mediator complex, a coactivator involved in the regulated

Reactivity:

bridge to convey information from gene-specific regulatory proteins to the basal RNA
polymerase II transcription machinery. Mediator is recruited to promoters by direct

interactions with regulatory proteins and serves as a scaffold for the assembly of a

Source: functional preinitiation complex with RNA polymerase II and the general transcription

functional preinitiation complex with RNA polymerase II and the general transcription factors. May be part of a complex containing NF2/merlin that participates in cellular

signaling to the actin cytoskeleton downstream of tyrosine kinase signaling pathways.

transcription of nearly all RNA polymerase II-dependent genes. Mediator functions as a

Isotype: Gene ID:

IgG MED28

Applications: Uniprot

ELISA Q9H204

Recommended dilutions: Synonyms:

Mediator of RNA polymerase II transcription subunit 28 (Endothelial-derived protein 1) (Mediator complex subunit 28) (Merlin and Grb2-interacting cytoskeletal protein)

(Magicin) (Tumor angiogenesis marker EG-1), MED28, EG1

Immunogen:

Recombinant Human Mediator of RNA polymerase II transcription subunit 28 protein (1-178AA).

Storage:

Preservative: 0.03% Proclin 300. Constituents: 50% Glycerol, 0.01M PBS, PH 7.4

Product	Images
---------	---------------

N/A N/A