
DDX19A Antibody

PACO42250




| Product Information | |
|---|--|
| Size: | Protein Background: |
| 50ug | ATP-dependent RNA helicase involved in mRNA export from the nucleus. Rather than unwinding RNA duplexes, DDX19 functions as a remodeler of ribonucleoprotein particles, whereby proteins bound to nuclear mRNA are dissociated and replaced by cytoplasmic mRNA binding proteins. |
| Reactivity: | |
| Human, Mouse | |
| Source: | Gene ID: |
| Rabbit | DDX19A |
| lsotype: | Uniprot |
| lgG | Q9NUU7 |
| Applications: | Synonyms: |
| ELISA, WB, IHC | ATP-dependent RNA helicase DDX19A (EC 3.6.4.13) (DDX19-like protein) (DEAD box protein 19A), DDX19A, DDX19L |
| Recommended dilutions: | Immunogen: |
| ELISA:1:2000-1:10000, WB:1:1000-1:5000, IHC:1:20-1:200 | Recombinant Human ATP-dependent RNA helicase DDX19A protein (1-478AA). |
| | Storage: |
| | Preservative: 0.03% Proclin 300. Constituents: 50% Glycerol, 0.01M PBS, PH 7.4 |



Western blot. All lanes: DDX19A antibody at 2μ g/ml + Mouse gonad tissue. Secondary. Goat polyclonal to rabbit IgG at 1/10000 dilution. Predicted band size: 54, 44 kDa. Observed band size: 54 kDa.

Immunohistochemistry of paraffin-embedded human prostate cancer using PACO42250 at dilution of 1:100.



Immunohistochemistry of paraffin-embedded human ovarian cancer using PACO42250 at dilution of 1:100.