

Product Information

Size:

50ul

Reactivity:

Human

Source:

Rabbit

Isotype:

IgG

Applications:

ELISA, IHC

Recommended dilutions:

ELISA:1:2000-1:10000, IHC:1:20-1:200

Protein Background:

Essential component of the COP9 signalosome complex (CSN), a complex involved in various cellular and developmental processes. The CSN complex is an essential regulator of the ubiquitin (Ubl) conjugation pathway by mediating the deneddylation of the cullin subunits of SCF-type E3 ligase complexes, leading to decrease the Ubl ligase activity of SCF-type complexes such as SCF, CSA or DDB2. The complex is also involved in phosphorylation of p53/TP53, c-jun/JUN, IkappaBalpha/NFKBIA, ITPK1 and IRF8/ICSBP, possibly via its association with CK2 and PKD kinases. CSN-dependent phosphorylation of TP53 and JUN promotes and protects degradation by the Ubl system, respectively. Involved in early stage of neuronal differentiation via its interaction with NIF3L1.

Gene ID:

COPS2

Uniprot

P61201

Synonyms:

COP9 signalosome complex subunit 2 (SGN2) (Signalosome subunit 2) (Alien homolog) (JAB1-containing signalosome subunit 2) (Thyroid receptor-interacting protein 15) (TR-interacting protein 15) (TRIP-15), COPS2, CSN2 TRIP15


Immunogen:

Recombinant Human COP9 signalosome complex subunit 2 protein (1-310AA).


Storage:

PBS with 0.02% sodium azide, 50% glycerol, pH7.3.

Product Images



Immunohistochemistry of paraffin-embedded human skin tissue using PACO43237 at dilution of 1:100.



Immunohistochemistry of paraffin-embedded human colon cancer using PACO43237 at dilution of 1:100.