## **ZMYND10 Antibody**



## PACO43808

Reactivity:

## **Product Information**

Size: **Protein Background:** 

50ul Required for motile ciliary function. Probably involved in axonemal assembly of inner

and outer dynein arms (IDA and ODA, respectively) for proper axoneme building for

cilia motility. May act by indirectly regulating transcription of dynein proteins.

Human, Mouse Gene ID:

Source: ZMYND10

Rabbit Uniprot

Isotype: O75800

lgG Synonyms:

**Applications:** Zinc finger MYND domain-containing protein 10 (Protein BLu), ZMYND10, BLU

ELISA, WB, IHC Immunogen:

**Recommended dilutions:** Recombinant Human Zinc finger MYND domain-containing protein 10 protein (201-


440AA). ELISA:1:2000-1:10000, WB:1:500-1:2000,

IHC:1:20-1:200


Storage:

PBS with 0.02% sodium azide, 50% glycerol, pH7.3.


## **Product Images**



Western blot. All lanes: ZMYND10 antibody at 8µg/ml. Lane 1: A549 whole cell lysate. Lane 2: Mouse kidney tissue. Lane 3: Mouse skeletal muscle tissue. Secondary. Goat polyclonal to rabbit IgG at 1/10000 dilution. Predicted band size: 51, 50 kDa. Observed band size: 51 kDa.



Immunohistochemistry of paraffin-embedded human lung cancer using PACO43808 at dilution of 1:100.



Immunohistochemistry of paraffin-embedded human testis tissue using PACO43808 at dilution of 1:100.