

PACO44552

Product Information

Size:

50ul

Reactivity:

Human

Source:

Rabbit

Isotype:

IgG

Applications:

ELISA, WB, IHC

Recommended dilutions:

ELISA:1:2000-1:10000, WB:1:1000-1:5000,
IHC:1:20-1:200

Protein Background:

Acts as a negative mediator of apoptosis and may play a role in tumor progression. Suppresses the TNF-mediated apoptosis by inhibiting caspase-8 activity but not the processing of procaspase-8, subsequently resulting in inhibition of BID cleavage and caspase-3 activation.

Gene ID:

TNFAIP8

Uniprot

O95379

Synonyms:

Tumor necrosis factor alpha-induced protein 8 (TNF alpha-induced protein 8) (Head and neck tumor and metastasis-related protein) (MDC-3.13) (NF-kappa-B-inducible DED-containing protein) (NDED) (SCC-S2) (TNF-induced protein GG2-1), TNFAIP8


Immunogen:

Recombinant Human Tumor necrosis factor alpha-induced protein 8 protein (1-198AA).


Storage:

PBS with 0.02% sodium azide, 50% glycerol, pH7.3.


Product Images



Western blot. All lanes: TNFAIP8 antibody at 2.32 μ g/ml. Lane 1: A431 whole cell lysate. Lane 2: PC-3 whole cell lysate. Secondary. Goat polyclonal to rabbit IgG at 1/10000 dilution. Predicted band size: 24, 23, 22, 25 kDa. Observed band size: 22 kDa.



Immunohistochemistry of paraffin-embedded human breast cancer using PACO44552 at dilution of 1:100.



Immunohistochemistry of paraffin-embedded human colon cancer using PACO44552 at dilution of 1:100.