

ZNF24 Antibody



PACO44855

Product Information

Size:

50ul

Reactivity:

Human

Source:

Rabbit

Isotype:

IgG

Applications:

ELISA, WB, IHC

Recommended dilutions:

ELISA:1:2000-1:10000, WB:1:1000-1:5000,
IHC:1:20-1:200

Protein Background:

Transcription factor required for myelination of differentiated oligodendrocytes. Required for the conversion of oligodendrocytes from the premyelinating to the myelinating state. In the developing central nervous system (CNS), involved in the maintenance in the progenitor stage by promoting the cell cycle. Specifically binds to the 5'-TCAT-3' DNA sequence. Has transcription repressor activity in vitro.

Gene ID:

ZNF24

Uniprot

P17028

Synonyms:

Zinc finger protein 24 (Retinoic acid, suppression protein A) (RSG-A) (Zinc finger and SCAN domain-containing protein 3) (Zinc finger protein 191) (Zinc finger protein KOX17), ZNF24, KOX17 ZNF191 ZSCAN3


Immunogen:

Recombinant Human Zinc finger protein 24 protein (169-368AA).


Storage:

PBS with 0.02% sodium azide, 50% glycerol, pH7.3.


Product Images



Western blot. All lanes: ZNF24 antibody at 2.88 μ g/ml. Lane 1: 293T whole cell lysate. Lane 2: Jurkat whole cell lysate. Lane 3: Hela whole cell lysate. Secondary. Goat polyclonal to rabbit IgG at 1/10000 dilution. Predicted band size: 43, 23 kDa. Observed band size: 43 kDa.



Immunohistochemistry of paraffin-embedded human kidney tissue using PACO44855 at dilution of 1:100.



Immunohistochemistry of paraffin-embedded human breast cancer using PACO44855 at dilution of 1:100.