MED25 Antibody, HRP conjugated

PACO46651



Product Information	
Size:	Protein Background:
50ug	Component of the Mediator complex, a coactivator involved in the regulated
Reactivity:	transcription of nearly all RNA polymerase II-dependent genes. Mediator functions as a bridge to convey information from gene-specific regulatory proteins to the basal RNA
Human	polymerase II transcription machinery. Mediator is recruited to promoters by direct interactions with regulatory proteins and serves as a scaffold for the assembly of a
Source:	functional preinitiation complex with RNA polymerase II and the general transcription
Rabbit	factors. Required for RARA/RXRA-mediated transcription.
lsotype:	Gene ID:
lgG	MED25
Applications:	Uniprot
ELISA	Q71SY5
Recommended dilutions:	Synonyms:
	Mediator of RNA polymerase II transcription subunit 25 (Activator interaction domain- containing protein 1) (Activator-recruited cofactor 92 kDa component) (ARC92) (Mediator complex subunit 25) (p78), MED25, acid, ARC92 PTOV2
	Immunogen:
	Recombinant Human Mediator of RNA polymerase II transcription subunit 25 protein (112-140AA).
	Storage:
	Preservative: 0.03% Proclin 300. Constituents: 50% Glycerol, 0.01M PBS, PH 7.4

N/A N/A