EZH1 Antibody, Biotin conjugated

PACO46977



Product Information	
Size:	Protein Background:
50ug	Polycomb group (PcG) protein. Catalytic subunit of the PRC2/EED-EZH1 complex, which
Reactivity:	methylates 'Lys-27' of histone H3, leading to transcriptional repression of the affected target gene. Able to mono-, di- and trimethylate 'Lys-27' of histone H3 to form
Human	H3K27me1, H3K27me2 and H3K27me3, respectively. Required for embryonic stem cell derivation and self-renewal, suggesting that it is involved in safeguarding embryonic
Source:	stem cell identity. Compared to EZH2-containing complexes, it is less abundant in
Rabbit	embryonic stem cells, has weak methyltransferase activity and plays a less critical role in forming H3K27me3, which is required for embryonic stem cell identity and proper
lsotype:	differentiation.
lgG	Gene ID:
Applications:	EZH1
ELISA	Uniprot
Recommended dilutions:	Q92800
	Synonyms:
	Histone-lysine N-methyltransferase EZH1 (EC 2.1.1.43) (ENX-2) (Enhancer of zeste homolog 1), EZH1, KIAA0388
	Immunogen:
	Recombinant Human Histone-lysine N-methyltransferase EZH1 protein (160-280AA).
	Storage:
	Preservative: 0.03% Proclin 300. Constituents: 50% Glycerol, 0.01M PBS, PH 7.4

N/A N/A