DNMT3B Antibody, HRP conjugated

PACO47159



Product Information	
Size:	Protein Background:
50ug	Required for genome-wide de novo methylation and is essential for the establishment
Reactivity:	of DNA methylation patterns during development. DNA methylation is coordinated with methylation of histones. May preferentially methylates nucleosomal DNA within
Human	the nucleosome core region. May function as transcriptional co-repressor by associating with CBX4 and independently of DNA methylation. Seems to be involved in
Source:	gene silencing. In association with DNMT1 and via the recruitment of CTCFL/BORIS,
Rabbit	involved in activation of BAG1 gene expression by modulating dimethylation of promoter histone H3 at H3K4 and H3K9. Isoforms 4 and 5 are probably not functional
lsotype:	due to the deletion of two conserved methyltransferase motifs. Function as transcriptional corepressor by associating with ZHX1.
lgG	Gene ID:
Applications:	DNMT3B
ELISA	Uniprot
Recommended dilutions:	Q9UBC3
	Synonyms:
	DNA (cytosine-5)-methyltransferase 3B (Dnmt3b) (EC 2.1.1.37) (DNA methyltransferase HsallIB) (DNA MTase HsallIB) (M. HsallIB), DNMT3B
	Immunogen:
	Recombinant Human DNA (cytosine-5)-methyltransferase 3B protein (1-100AA).
	Storage:

Preservative: 0.03% Proclin 300. Constituents: 50% Glycerol, 0.01M PBS, PH 7.4

N/A N/A