TTI1 Antibody



PACO47494

Reactivity:

Rabbit

Product Information

Size: Protein Background:

50ug Regulator of the DNA damage response (DDR). Part of the TTT complex that is required

to stabilize protein levels of the phosphatidylinositol 3-kinase-related protein kinase (PIKK) family proteins. The TTT complex is involved in the cellular resistance to DNA

Human damage stresses, like ionizing radiation (IR), ultraviolet (UV) and mitomycin C (MMC).

Together with the TTT complex and HSP90 may participate in the proper folding of

Source: newly synthesized PIKKs. Promotes assembly, stabilizes and maintains the activity of

mTORC1 and mTORC2 complexes, which regulate cell growth and survival in response

to nutrient and hormonal signals.

Isotype: Gene ID:

IgG TTI1

Applications: Uniprot

ELISA, IHC, IF O43156

Recommended dilutions: Synonyms:

ELISA:1:2000-1:10000, IHC:1:20-1:200, TELO2-interacting protein 1 homolog (Protein SMG10), TTI1, KIAA0406 SMG10 IF:1:50-1:200


Immunogen:

Recombinant Human TELO2-interacting protein 1 homolog protein (692-858AA).


Storage:

Preservative: 0.03% Proclin 300. Constituents: 50% Glycerol, 0.01M PBS, PH 7.4


Product Images



Immunohistochemistry of paraffin-embedded human heart tissue using PACO47494 at dilution of 1:100.



Immunofluorescent analysis of Hela cells using PACO47494 at dilution of 1:100 and Alexa Fluor 488-congugated AffiniPure Goat Anti-Rabbit IgG(H+L).



Immunohistochemistry of paraffin-embedded human kidney tissue using PACO47494 at dilution of 1:100.