## **SLC33A1 Antibody**



## PACO49122

**Product Information** 

Size: **Protein Background:** 

50ug Probable acetyl-CoA transporter necessary for O-acetylation of gangliosides. Negatively

regulates BMP signaling.

Reactivity: Gene ID:

Human, Mouse SLC33A1

Source:

Uniprot Rabbit

O00400 Isotype:

Synonyms: lgG

Acetyl-coenzyme A transporter 1 (AT-1) (Acetyl-CoA transporter 1) (Solute carrier family **Applications:** 

33 member 1), SLC33A1, ACATN AT1

ELISA, WB, IHC Immunogen:

**Recommended dilutions:** Recombinant Human Acetyl-coenzyme A transporter 1 protein (1-74AA).


ELISA:1:2000-1:10000, WB:1:1000-1:5000,

IHC:1:20-1:200


Storage:

Preservative: 0.03% Proclin 300. Constituents: 50% Glycerol, 0.01M PBS, PH 7.4


## **Product Images**



Immunohistochemistry of paraffin-embedded human thyroid tissue using PACO49122 at dilution of 1:100.



Western Blot. Positive WB detected in: A549 whole cell lysate, HepG2 whole cell lysate, Mouse liver tissue, Mouse kidney tissue, Mouse brain tissue. All lanes: SLC33A1 antibody at 2.5µg/ml. Secondary. Goat polyclonal to rabbit lgG at 1/50000 dilution. Predicted band size: 61 kDa. Observed band size: 50 kDa.



Immunohistochemistry of paraffin-embedded human kidney tissue using PACO49122 at dilution of 1:100.