## EIF3J Antibody, HRP conjugated

## PACO49551



Product Information	
Size:	Protein Background:
50ug	Component of the eukaryotic translation initiation factor 3 (eIF-3) complex, which is
Reactivity:	required for several steps in the initiation of protein synthesis. The eIF-3 complex associates with the 40S ribosome and facilitates the recruitment of eIF-1, eIF-1A, eIF-2:
Human	GTP: methionyl-tRNAi and eIF-5 to form the 43S preinitiation complex (43S PIC). The eIF-3 complex stimulates mRNA recruitment to the 43S PIC and scanning of the mRNA
Source:	for AUG recognition. The eIF-3 complex is also required for disassembly and recycling
Rabbit	of post-termination ribosomal complexes and subsequently prevents premature joining of the 40S and 60S ribosomal subunits prior to initiation. This subunit binds directly
lsotype:	within the mRNA entry channel of the 40S ribosome to the aminoacyl (A) site. It may regulate the interaction between the 43S PIC and mRNA.
lgG	Gene ID:
Applications:	EIF3J
ELISA	Uniprot
Recommended dilutions:	O75822
	Synonyms:
	Eukaryotic translation initiation factor 3 subunit J (eIF3j) (Eukaryotic translation initiation factor 3 subunit 1) (eIF-3-alpha) (eIF3 p35), EIF3J, EIF3S1
	Immunogen:
	Recombinant Human Eukaryotic translation initiation factor 3 subunit J protein (5- 115AA).
	Storage:

Preservative: 0.03% Proclin 300. Constituents: 50% Glycerol, 0.01M PBS, pH 7.4

N/A N/A