WNT5A Antibody, FITC conjugated



PACO50096

Reactivity:

Isotype:

lgG

Product Information

Size: Protein Background:

50ug Ligand for members of the frizzled family of seven transmembrane receptors. Can activate or inhibit canonical Wnt signaling, depending on receptor context. In the

presence of FZD4, activates beta-catenin signaling. In the presence of ROR2, inhibits the

Human canonical Wnt pathway by promoting beta-catenin degradation through a GSK3-independent pathway which involves down-regulation of beta-catenin-induced

Source: reporter gene expression. Suppression of the canonical pathway allows chondrogenesis

Rabbit to occur and inhibits tumor formation. Stimulates cell migration. Decreases

proliferation, migration, invasiveness and clonogenicity of carcinoma cells and may act

as a tumor suppressor. Mediates motility of melanoma cells. Required during embryogenesis for extension of the primary anterior-posterior axis and for outgrowth

of limbs and the genital tubercle. Inhibits type II collagen expression in chondrocytes.

Applications: Gene ID:

ELISA WNT5A

Recommended dilutions: Uniprot

P41221

Synonyms:

Protein Wnt-5a, WNT5A

Immunogen:

Recombinant Human Protein Wnt-5a protein (62-380AA).

Storage:

Preservative: 0.03% Proclin 300. Constituents: 50% Glycerol, 0.01M PBS, pH 7.4

Product	Images
---------	---------------

N/A N/A