SLAMF1 Antibody, Biotin conjugated



PACO50493

Reactivity:

Human

Product Information

Recommended dilutions:

Size: Protein Background:

50ug Self-ligand receptor of the signaling lymphocytic activation molecule (SLAM) family.

SLAM receptors triggered by homo- or heterotypic cell-cell interactions are modulating

the activation and differentiation of a wide variety of immune cells and thus are involved in the regulation and interconnection of both innate and adaptive immune

response. Activities are controlled by presence or absence of small cytoplasmic adapter proteins, SH2D1A/SAP and/or SH2D1B/EAT-2. SLAMF1-induced signal-transduction

Rabbit events in T-lymphocytes are different from those in B-cells. Two modes of SLAMF1 signaling seem to exist: one depending on SH2D1A (and perhaps SH2D1B) and another

Isotype: in which protein-tyrosine phosphatase 2C (PTPN11)-dependent signal transduction

operates.

lgG **Gene ID:**

Applications: SLAMF1

ELISA Uniprot

Q13291

Synonyms:

Signaling lymphocytic activation molecule (CDw150) (IPO-3) (SLAM family member 1) (CD antigen CD150), SLAMF1, SLAM

Immunogen:

Recombinant Human Signaling lymphocytic activation molecule protein (69-204AA).

Storage:

Preservative: 0.03% Proclin 300. Constituents: 50% Glycerol, 0.01M PBS, pH 7.4

Product	Images
---------	---------------

N/A N/A