

PACO50582

Product Information

Size:

50ug

Reactivity:

Human, Rat

Source:

Rabbit

Isotype:

IgG

Applications:

ELISA, WB, IHC, IF, IP

Recommended dilutions:

ELISA:1:2000-1:10000, WB:1:500-1:5000,
IHC:1:20-1:200, IF:1:50-1:200, IP:1:200-
1:2000,

Protein Background:

The nuclear form acts as a tumor suppressor by preventing ubiquitination and degradation of isoform p19ARF/ARF of CDKN2A by TRIP12: acts by interacting with TRIP12, leading to disrupt interaction between TRIP12 and isoform p19ARF/ARF of CDKN2A. Adapter molecule for TNFRSF1A/TNFR1 that specifically associates with the cytoplasmic domain of activated TNFRSF1A/TNFR1 mediating its interaction with FADD. Overexpression of TRADD leads to two major TNF-induced responses, apoptosis and activation of NF-kappa-B.

Gene ID:

TRADD

Uniprot

Q15628

Synonyms:

Tumor necrosis factor receptor type 1-associated DEATH domain protein (TNFR1-associated DEATH domain protein) (TNFRSF1A-associated via death domain), TRADD


Immunogen:

Recombinant Human Tumor necrosis factor receptor type 1-associated DEATH domain protein (1-312AA).


Storage:

Preservative: 0.03% Proclin 300. Constituents: 50% Glycerol, 0.01M PBS, pH 7.4


Product Images



Western Blot. Positive WB detected in: K562 whole cell lysate, Rat kidney tissue. All lanes: TRADD antibody at 4 μ g/ml. Secondary: Goat polyclonal to rabbit IgG at 1/50000 dilution. Predicted band size: 35, 28 kDa. Observed band size: 35 kDa.



Immunohistochemistry of paraffin-embedded human tonsil tissue using PACO50582 at dilution of 1:100.



Immunohistochemistry of paraffin-embedded human liver tissue using PACO50582 at dilution of 1:100.