ATG4C Antibody, HRP conjugated

PACO50827



Product Information	
Size:	Protein Background:
50ug	Cysteine protease required for the cytoplasm to vacuole transport (Cvt) and autophagy.
Reactivity:	Is not essential for autophagy development under normal conditions but is required for a proper autophagic response under stressful conditions such as prolonged starvation.
Human	Cleaves the C-terminal amino acid, of ATG8 family proteins MAP1LC3 and GABARAPL2, to reveal a C-terminal glycine. Exposure of the glycine at the C-terminus is essential for
Source:	ATG8 proteins conjugation to phosphatidylethanolamine (PE) and insertion to
Rabbit	membranes, which is necessary for autophagy. Has also an activity of delipidating enzyme for the PE-conjugated forms.
lsotype:	Gene ID:
lgG	ATG4C
Applications:	Uniprot
ELISA	Q96DT6
Recommended dilutions:	Synonyms:
	Cysteine protease ATG4C (EC 3.4.22) (AUT-like 3 cysteine endopeptidase) (Autophagin- 3) (Autophagy-related cysteine endopeptidase 3) (Autophagy-related protein 4 homolog C), ATG4C, APG4C AUTL1 AUTL3
	Immunogen:
	Recombinant Human Cysteine protease ATG4C protein (419-448AA).
	Storage:
	Preservative: 0.03% Proclin 300. Constituents: 50% Glycerol, 0.01M PBS, pH 7.4

N/A N/A