

PACO51030

---

## Product Information

**Size:**

50ug

**Reactivity:**

Bacillus sp.

**Source:**

Rabbit

**Isotype:**

IgG

**Applications:**

ELISA, WB

**Recommended dilutions:**

ELISA:1:2000-1:10000, WB:1:500-1:5000

**Protein Background:**

Catalyzes the hydrolysis of oligosaccharides with unsaturated glucuronyl residues at the non-reducing terminal, to a sugar or an amino sugar, and an unsaturated D-glucuronic acid, (GlcA), which is nonenzymatically converted immediately to alpha-keto acid, .

**Gene ID:**

ugl

**Uniprot**

Q9RC92

**Synonyms:**

Unsaturated glucuronyl hydrolase (UGL) (EC 3.2.1.179) (Glycosaminoglycan hydrolase) (Glycuronidase) (Unsaturated uronic acid, hydrolase), ugl

**Immunogen:**


Recombinant Bacillus sp. Unsaturated glucuronyl hydrolase protein (1-377AA).

**Storage:**

Preservative: 0.03% Proclin 300. Constituents: 50% Glycerol, 0.01M PBS, pH 7.4

## Product Images

---



Western Blot. Positive WB detected in Recombinant protein. All lanes: ugi antibody at 2.8 $\mu$ g/ml. Secondary. Goat polyclonal to rabbit IgG at 1/50000 dilution. predicted band size: 43 kDa. observed band size: 43 kDa.