## TFB2M Antibody, FITC conjugated



## PACO51040

Source:

## **Product Information**

Size: Protein Background:

50ug S-adenosyl-L-methionine-dependent methyltransferase which specifically dimethylates

**Reactivity:** mitochondrial 12S rRNA at the conserved stem loop. Also required for basal transcription of mitochondrial DNA, probably via its interaction with POLRMT and

Human TFAM. Stimulates transcription independently of the methyltransferase activity.

Compared to TFB1M, it activates transcription of mitochondrial DNA more efficiently,

while it has less methyltransferase activity.

Rabbit Gene ID:

Isotype: TFB2M

lgG Uniprot

Applications: Q9H5Q4

ELISA Synonyms:

**Recommended dilutions:** Dimethyladenosine transferase 2, mitochondrial (EC 2.1.1) (Hepatitis C virus NS5A-

transactivated protein 5) (HCV NS5A-transactivated protein 5) (Mitochondrial 12S rRNA dimethylase 2) (Mitochondrial transcription factor B2) (h-mtTFB) (h-mtTFB2) (hTFB2M) (mtTFB2) (S-adenosylmethionine-6-N', N'-adenosyl(rRNA) dimethyltransferase 2),

TFB2M, NS5ATP5

Immunogen:

Recombinant Human Dimethyladenosine transferase 2, mitochondrial protein (30-

108AA).

Storage:

Preservative: 0.03% Proclin 300. Constituents: 50% Glycerol, 0.01M PBS, pH 7.4

Product	<b>Images</b>
---------	---------------

N/A N/A