## **CELF2 Antibody**

# **Assay**Genie

#### PACO51458

#### **Product Information**

Size: **Protein Background:** 

50ug

**Applications:** 

IHC:1:500-1:1000, IF:1:50-1:200

Reactivity: Gene ID:

CELF2 Human

Source: Uniprot

O95319 Rabbit

Isotype: **Synonyms:** 

lgG CUGBP Elav-like family member 2 (CELF-2) (Bruno-like protein 3) (CUG triplet repeat

> RNA-binding protein 2) (CUG-BP2) (CUG-BP- and ETR-3-like factor 2) (ELAV-type RNAbinding protein 3) (ETR-3) (Neuroblastoma apoptosis-related RNA-binding protein)

(hNAPOR) (RNA-binding protein BRUNOL-3), CELF2, BRUNOL3 CUGBP2 ETR3 NAPOR ELISA, WB, IHC, IF


Immunogen: **Recommended dilutions:** 

Recombinant Human CUGBP Elav-like family member 2 protein (291-381AA). ELISA:1:2000-1:10000, WB:1:500-1:5000,


Storage:


Preservative: 0.03% Proclin 300. Constituents: 50% Glycerol, 0.01M PBS, pH 7.4

### **Product Images**



Western Blot. Positive WB detected in: K562 whole cell lysate, A549 whole cell lysate. All lanes: CELF2 antibody at  $3\mu g/ml$ . Secondary. Goat polyclonal to rabbit lgG at 1/50000 dilution. Predicted band size: 55, 53, 56 kDa. Observed band size: 55 kDa.





IHC image of PACO51458 diluted at 1:600 and staining in paraffinembedded human brain tissue performed on a Leica BondTM system. After dewaxing and hydration, antigen retrieval was mediated by high pressure in a citrate buffer (pH 6.0). Section was blocked with 10% normal goat serum 30min at RT. Then primary antibody (1% BSA) was incubated at 4°C overnight. The primary is detected by a biotinylated secondary antibody and visualized using an HRP conjugated SP system.

IHC image of PACO51458 diluted at 1:600 and staining in paraffinembedded human tonsil tissue performed on a Leica BondTM system. After dewaxing and hydration, antigen retrieval was mediated by high pressure in a citrate buffer (pH 6.0). Section was blocked with 10% normal goat serum 30min at RT. Then primary antibody (1% BSA) was incubated at 4°C overnight. The primary is detected by a biotinylated secondary antibody and visualized using an HRP conjugated SP system.