

PACO53514

Product Information

Size:

50ug

Reactivity:

Human

Source:

Rabbit

Isotype:

IgG

Applications:

ELISA, WB, IF

Recommended dilutions:

ELISA:1:2000-1:10000, WB:1:500-1:5000,
IF:1:50-1:200

Protein Background:

Plays an important role in deacetylating mercapturic acid, in kidney proximal tubules. Also acts on N-acetyl-aromatic amino acid, .

Gene ID:

ACY3

Uniprot

Q96HD9

Synonyms:

N-acyl-aromatic-L-amino acid, amidohydrolase (carboxylate-forming) (EC 3.5.1.114) (Acylase III) (Aminoacylase-3) (ACY-3) (Aspartoacylase-2) (Hepatitis C virus core-binding protein 1) (HCBP1) (HCV core-binding protein 1), ACY3, ASPA2


Immunogen:

Recombinant Human N-acyl-aromatic-L-amino acid, amidohydrolase (carboxylate-forming) protein (5-319AA).


Storage:

Preservative: 0.03% Proclin 300. Constituents: 50% Glycerol, 0.01M PBS, pH 7.4

Product Images



Western Blot. Positive WB detected in: MCF-7 whole cell lysate. All lanes: ACY3 antibody at 3 μ g/ml. Secondary. Goat polyclonal to rabbit IgG at 1/50000 dilution. Predicted band size: 36 kDa. Observed band size: 36 kDa.



Immunofluorescent analysis of HepG2 cells using PACO53514 at dilution of 1:100 and Alexa Fluor 488-conjugated AffiniPure Goat Anti-Rabbit IgG(H+L).