WNK2 Antibody, Biotin conjugated

PACO54233



Product Information	
Size:	Protein Background:
50ug	Serine/threonine kinase which plays an important role in the regulation of electrolyte
Reactivity:	homeostasis, cell signaling, survival, and proliferation. Acts as an activator and inhibitor of sodium-coupled chloride cotransporters and potassium-coupled chloride
Human	cotransporters respectively. Activates SLC12A2, SCNN1A, SCNN1B, SCNN1D and SGK1 and inhibits SLC12A5. Negatively regulates the EGF-induced activation of the
Source:	ERK/MAPK-pathway and the downstream cell cycle progression. Affects
Rabbit	MAPK3/MAPK1 activity by modulating the activity of MAP2K1 and this modulation depends on phosphorylation of MAP2K1 by PAK1. WNK2 acts by interfering with the
lsotype:	activity of PAK1 by controlling the balance of the activity of upstream regulators of PAK1 activity, RHOA and RAC1, which display reciprocal activity.
lgG	Gene ID:
Applications:	WNK2
ELISA	Uniprot
Recommended dilutions:	Q9Y3S1
	Synonyms:
	Serine/threonine-protein kinase WNK2 (EC 2.7.11.1) (Antigen NY-CO-43) (Protein kinase lysine-deficient 2) (Protein kinase with no lysine 2) (Serologically defined colon cancer antigen 43), WNK2, KIAA1760 PRKWNK2 SDCCAG43
	Immunogen:
	Recombinant Human Serine/threonine-protein kinase WNK2 protein (1177-1464AA).
	Storage:

Preservative: 0.03% Proclin 300. Constituents: 50% Glycerol, 0.01M PBS, pH 7.4

N/A N/A