PRKCB Antibody, HRP conjugated



PACO55079

lgG

Product Information

Recommended dilutions:

Size: Protein Background:

50ug Calcium-activated, phospholipid- and diacylglycerol (DAG)-dependent

Reactivity:of the B-cell receptor (BCR) signalosome, oxidative stress-induced apoptosis, androgen receptor-dependent transcription regulation, insulin signaling and endothelial cells

proliferation. Plays a key role in B-cell activation by regulating BCR-induced NF-kappa-B

Source: activation. Mediates the activation of the canonical NF-kappa-B pathway (NFKB1) by

Rabbit direct phosphorylation of CARD11/CARMA1 at 'Ser-559', 'Ser-644' and 'Ser-652'.

Phosphorylation induces CARD11/CARMA1 association with lipid rafts and recruitment

Isotype: of the BCL10-MALT1 complex as well as MAP3K7/TAK1, which then activates IKK

complex, resulting in nuclear translocation and activation of NFKB1.

Gene ID:

Applications:

PRKCB

ELISA Uniprot

P05771

Synonyms:

Protein kinase C beta type (PKC-B) (PKC-beta) (EC 2.7.11.13), PRKCB, PKCB PRKCB1

serine/threonine-protein kinase involved in various cellular processes such as regulation

Immunogen:

Recombinant Human Protein kinase C β type protein (517-643AA).

Storage:

Preservative: 0.03% Proclin 300. Constituents: 50% Glycerol, 0.01M PBS, pH 7.4

Product	Images
---------	---------------

N/A N/A