GDF10 Antibody



PACO55518

Reactivity:

Product Information

Size: Protein Background:

50ug Growth factor involved in osteogenesis and adipogenesis. Plays an inhibitory role in the

process of osteoblast differentiation via SMAD2/3 pathway. Plays an inhibitory role in

the process of adipogenesis.

Human, Rat Gene ID:

Source: GDF10

Rabbit **Uniprot**

Isotype: P55107

lgG Synonyms:

Applications: Growth/differentiation factor 10 (GDF-10) (Bone morphogenetic protein 3B) (BMP-3B)

ELISA, WB, IHC (Bone-inducing protein) (BIP), GDF10, BMP3B

Recommended dilutions:

ELISA:1:2000-1:10000, WB:1:500-1:5000,


IHC:1:20-1:200

Recombinant Human Growth/differentiation factor 10 protein (369-478AA).


Storage:

Preservative: 0.03% Proclin 300. Constituents: 50% Glycerol, 0.01M PBS, pH 7.4

Product Images



Western Blot. Positive WB detected in: Rat brain tissue. All lanes: GDF10 antibody at $4.5\mu g/ml$. Secondary. Goat polyclonal to rabbit IgG at 1/50000 dilution. Predicted band size: 54 kDa. Observed band size: 54 kDa.



IHC image of PACO55518 diluted at 1:200 and staining in paraffinembedded human testis tissue performed on a Leica BondTM system. After dewaxing and hydration, antigen retrieval was mediated by high pressure in a citrate buffer (pH 6.0). Section was blocked with 10% normal goat serum 30min at RT. Then primary antibody (1% BSA) was incubated at 4°C overnight. The primary is detected by a biotinylated secondary antibody and visualized using an HRP conjugated SP system.

IHC image of PACO55518 diluted at 1:200 and staining in paraffinembedded human prostate tissue performed on a Leica BondTM system. After dewaxing and hydration, antigen retrieval was mediated by high pressure in a citrate buffer (pH 6.0). Section was blocked with 10% normal goat serum 30min at RT. Then primary antibody (1% BSA) was incubated at 4°C overnight. The primary is detected by a biotinylated secondary antibody and visualized using an HRP conjugated SP system.